

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE DEC 1974		2. REPORT TYPE		3. DATES COVERED 00-00-1974 to 00-00-1974	
4. TITLE AND SUBTITLE Occurrence of Two Types of Gynandromorphism in a Sibling Series of Aedes (Stegomyia) Craggi (Barraud) (Diptera: Culicidae)				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Smithsonian Institution, Medical Entomology Project, Department of Entomology, Washington, DC, 20560				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT see report					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

OCCURRENCE OF TWO TYPES OF GYNANDROMORPHISM IN A SIBLING SERIES OF *Aedes (Stegomyia) craggi* (BARRAUD) (DIPTERA: CULICIDAE)¹

YIAU-MIN HUANG

Medical Entomology Project, Department of Entomology, Smithsonian Institution,
Washington, D.C. 20560

ABSTRACT. Anterior-posterior and bilateral gynanders of *Aedes (Stegomyia) craggi* (Barraud) were encountered in a sibling series of this species. The occurrence of two types of gynandromorphism

within such a series of mosquitoes has not been previously reported. Reports of mosquito gynandromorphs subsequent to, or not included in the tabulation of Brust (1966) are summarized.

Two gynandromorphs of *Aedes (Stegomyia) craggi* (Barraud) were found among the *Stegomyia* mosquitoes submitted by the SEATO Medical Research Laboratory, Bangkok. Both specimens were reared from eggs obtained from a wild-caught female biting man in a forest at Chiang Mai, Thailand. It is noteworthy that these 2 gynandromorphs are of 2 distinct types known as an anterior-posterior gynander and a bilateral gynander. Accompanying these 2 gynanders are another 12 specimens (7 males, 5 females) which were derived from the same female (mother). These are normal siblings. As far as it can be determined, this is the first time that 2 types of gynandromorphism have been reported from the

same sibling series. Craig and Hickey (1969:102) report 4 gynanders in one sibship of 16 individuals but do not indicate if 2 or more types were represented. The two gynanders are as follows:

(1). Anterior-posterior gynander, specimen No. (1)-2 with associated terminalia on slide (SEAMP 345, 73/302). This specimen has antennae, male; palpi, male; fore- and midtarsal claws unequal, male; hindtarsomere 4 with basal 5/6 white and tarsomeres 3, 5 dark, female; abdomen and genitalia, female, normal, all three spermathecae present.

(2). Bilateral gynander, specimen No. (1)-5 with associated terminalia on slide (SEAMP 345, 73/301). This specimen has right antenna, male; right palpus, male; right fore- and midtarsal claws unequal, male; right hindtarsomeres 3-5 dark, male; left antenna, female; left palpus, female; left fore- and midtarsal claws equal, female; left hindtarsomere 4 with

¹ This work was supported by Research Contracts DA-49-193-MD-2672 and DAMD-17-74-C-4086 from the U.S. Army Medical Research Development Command, Office of the Surgeon General, Washington, D.C.

basal 5/6 white and tarsomeres 3, 5 dark, female; genitalia, male, normal.

Brust (1966) gives a detailed review of gynandromorphism in mosquitoes and cites much of the literature. Since then a number of gynandromorphs have been reported. The information is generally scattered throughout the literature. At present, 34 species belonging to 10 genera have been described. Table 1 summarizes published records of gynandromorphs which were omitted from the tabulation in Brust (1966) or have been reported subsequently.

ACKNOWLEDGMENTS. I am grateful to Dr. Ronald A. Ward for review of the manuscript and for his valuable comments; to Mr. E. L. Peyton for his sup-

port and to Mr. R. H. Hochman for his help in searching the literature. I also wish to express my gratitude to Dr. George B. Craig, Jr., Department of Biology, University of Notre Dame, Notre Dame, Indiana, for information concerning gynandromorphism in mosquitoes and for his kindness during my visit to the Vector Biology Laboratory, University of Notre Dame.

Literature Cited

- Aslamkhan, M. and R. H. Baker. 1969. Gynandromorphism in *Culex tritaeniorhynchus*. Mosq. News 29(1): 127-132.
 Bengé, M. K. 1970. A gynandromorph of *Culiseta inornata* (Williston). Mosq. News 30(2):268.
 Blakeslee, T. E. and P. T. Rigby. 1965. Gyn-

TABLE 1. Gynandromorphs in mosquitoes (not tabulated in Brust, 1966).

Genus	Subgenus	Species	Reference
<i>Aedes</i>	<i>(Aedes)</i>	<i>cinereus</i> Meigen	Brust (1966)
	<i>(Aedimorphus)</i>	<i>dentatus</i> (Theobald)	Van Someren (1969)
		<i>vexans</i> (Meigen)	Minson (1969); Horsfall <i>et al.</i> (1973)
	<i>(Finlaya)</i>	<i>togoi</i> (Theobald)	Chellappah (1965)
	<i>(Ochlerotatus)</i>	<i>dorsalis</i> (Meigen)	Blakeslee, Rigby and Bomotti (1966)
	<i>(Protomacleaya)</i>	<i>excrucians</i> (Walker)	Brust (1966)
		<i>triseriatus</i> (Say)	Ezenwa and Venard (1973)
		<i>pembaensis</i> Theobald	Paterson and Worth (1961)
		<i>aegypti</i> (Linnaeus)	Craig and Hickey (1967)
	<i>(Skusea)</i>	<i>albopictus</i> (Skuse)	Craig and Hickey (1967)
	<i>(Stegomyia)</i>	<i>giveni</i> Edwards	Colless (1958) ²
	<i>(Armigeres)</i>	<i>perturbans</i> (Walker)	Pinger (1972) ³
<i>Armigeres</i>	<i>(Coquillettidia)</i>	<i>erythrothorax</i> Dyar	Blakeslee and Rigby (1965)
<i>Coquillettidia</i>	<i>(Culex)</i>	<i>nigripalpus</i> Theobald	Meadows (1966); Taylor, Meadows and Branch (1966)
<i>Culex</i>		<i>quinquefasciatus</i> Say	Meadows (1966); Seal (1966); Taylor, Meadows and Branch (1966)
		<i>salinarius</i> Coquillett	Meadows (1966); Taylor, Meadows and Branch (1966)
		<i>tarsalis</i> Coquillett	Harmston (1965, 1971); Rigby (1966); Taylor, Meadows and Branch (1966); Rosay (1968); Mitchell and Hughes (1969)
		<i>tritaeniorhynchus</i> Giles	Aslamkhan and Baker (1969)
<i>Culiseta</i>	<i>(Culiciomyia)</i>	<i>cinereus</i> Theobald	Van Someren (1969)
	<i>(Culiseta)</i>	<i>inornata</i> (Williston)	Benge (1970)
	<i>(Climacura)</i>	<i>novaezealandiae</i> Pillai	Dobrotworsky (1972)
<i>Trichoprosopon</i>	<i>(Trichoprosopon)</i>	<i>digitatum</i> (Rondani)	Lee (1967)

² This reference is not available. The information is from Chellappah (1965).

³ Pinger (1972) reported it as an intersex. It is a gynandromorph (G. B. Craig, Jr., personal communication).

- andromorphism in *Culex erythrorhox* (Dyar). Mosq. News 25(2):219.
- Blakeslee, T. E., P. T. Rigby and G. Bomotti. 1966. An *Aedes dorsalis* (Meigen) gynandromorph. Calif. Vector Views 13(5):41.
- Brust, R. A. 1966. Gynandromorphs and intersexes in mosquitoes (Diptera: Culicidae). Can. J. Zool. 44:911-921.
- Chellappah, W. T. 1965. A gynandromorph of *Aedes* (F.) *togoi* (Theobald). Mosq. News 25(4):483-484.
- Colless, D. H. 1958. Notes on the Culicine mosquitoes of Singapore. Ph.D. thesis, Univ. Singapore.
- Craig, G. B., Jr. and W. A. Hickey. 1967. Genetics of *Aedes aegypti*. P. 67-131. In Wright, J. W. & R. Pal (eds.) Genetics of Insect Vectors of Disease. Elsevier Publishing Company, Amsterdam.
- Dobrotworsky, N. V. 1972. A gynandromorph of *Culiseta novaeseelandiae* Pillai (Diptera: Culicidae). J. Aust. Ent. Soc. 11(2):150-151.
- Ezenwa, A. O. and C. E. Venard. 1973. Gynandromorphism in *Aedes triseriatus* (Say) J. Med. Ent. 10(1):52.
- Harmston, F. G. 1965. Gynandromorph of *Culex tarsalis* Coquillett from Colorado. Mosq. News 25(4):488-489.
- . 1971. Two mosaic gynandromorphs of *Culex tarsalis* Coquillett from Texas. Mosq. News 31:222-223.
- Horsfall, W. R., H. W. Fowler, Jr., L. J. Moretti and J. R. Larsen. 1973. Bionomics and embryology of the inland floodwater mosquito *Aedes vexans*. Univ. Ill. Press, Urbana. 211 pp.
- Lee, V. H. 1967. Gynandromorphism in the Sabethine, *Trichoprosopon digitatum* (Rondani). Mosq. News 27(3):426-427.
- Meadows, K. E. 1966. Gynandromorphism in *Culex* (Linnaeus) mosquitoes, Tampa Bay Area, Florida—1965. Mosq. News 26(4):587-589.
- Minson, K. L. 1969. An *Aedes vexans* gynandromorph. Mosq. News 29(1):135.
- Mitchell, C. J. and T. B. Hughes. 1969. A bipolar differentiated gynandromorph of *Culex tarsalis* Coquillett from Texas. J. Med. Ent. 6:78.
- Paterson, H. E. and C. B. Worth. 1961. Gynandromorphism in an African mosquito (Diptera: Culicidae) J. Ent. Soc. S. Africa 24(1):214-215.
- Pinger, R. R., Jr. 1972. An intersex of *Mansonia perturbans*. Mosq. News 32(4):633-634.
- Rigby, P. T. 1966. Gynandromorphism in *Culex tarsalis* Coquillett. Mosq. News 26:584.
- Rosay, B. 1968. An autogenous gynandromorph of *Culex tarsalis* Coq. Mosq. News 28(1):120.
- Seal, C. W. 1966. A gynandromorph of *Culex pipiens quinquefasciatus* (Say). Mosq. News 26(4):586-587.
- Someran, E. C. C. Van. 1969. Some interesting mosquitoes from Kenya. Mosq. Syst. Newsl. 1(2):28.
- Taylor, D., K. Meadows and N. Branch. 1966. Gynandromorphism in *Culex* (Linnaeus) mosquitoes collected in the Tampa Bay Area 1962 through 1964. Mosq. News 26(1):8-10.